

Links between connections, parallel transports and transports along paths

Bozhidar Z. Iliev * [†] [‡]

Short title: Connections and (parallel) transports along paths

Last update: → February 29, 2008

*Report presented at the International Workshop
“Advanced Geometric Methods in Physics”
Florence, Italy, 14 – 18 April, 2005*

<http://www.arXiv.org> e-Print archive No.: math.DG/0504010

BOZHO[®] TM

Subject Classes:

Differential geometry

2000 MSC numbers:

53C05, 53C99

55R99, 58A30

2003 PACS numbers:

02.40.Ma, 02.40.Vh

02.40.-k, 04.20.Cv

Key- Words:

Connections on bundles, Parallel transport

Axiomatically defined parallel transport, Transports along paths in bundles

*Laboratory of Mathematical Modeling in Physics, Institute for Nuclear Research and Nuclear Energy, Bulgarian Academy of Sciences, Boul. Tzarigradsko chaussée 72, 1784 Sofia, Bulgaria

[†]E-mail address: bozho@inrne.bas.bg

[‡]URL: <http://theo.inrne.bas.bg/~bozho/>

Contents

1	Introduction	1
2	Connections and the generated by them parallel transports (review)	3
3	Transports along paths in fibre bundles (review)	5
4	A connection generated by transport along paths	6
5	Links between parallel transports and transports along paths	8
6	Relations between connections and (parallel) transports along paths	10
7	Recapitulation	11
8	The case of topological bundles	12
9	The case of vector bundles	15
10	Conclusion	17
	References	18
	This article ends at page	21

Abstract

The axiomatic approach to parallel transport theory is partially discussed. Bijective correspondences between the sets of connections, (axiomatically defined) parallel transports and transports along paths satisfying some additional conditions are constructed. In particular, the equivalence between the concepts “connection” and “(axiomatically defined) parallel transport” is established. Some results which are specific for topological and vector bundles are presented.

1. Introduction

From different view-points, the connection theory can be found in many works, like [1–21]. As pointed in these and many other references, the concept of a parallel transport is defined on the base of the one of a connection. The opposite approach, i.e. the definition of a connection on the ground of an axiomatically defined one of a parallel transport, is considered in [22–32]. The major classical results on axiomatization of parallel transport theory are presented in the table below

Year	Person	Result and original reference
1917	T. Levi-Civita	Definition of a parallel transport of a vector in Riemannian geometry. [33]
1949	Willi Rinow	An axiomatic definition of parallel transport in tangent bundle is introduced in unpublished lectures at Humboldt university. (See [25] and [32, p. 46].)
1964	Ü.G. Lumiste	Definition of a connection in principal bundle (with homogeneous fibres) as a parallel transport along canonical paths $\alpha: [0, 1] \rightarrow M$ in its base M . The parallel transport is defined as a mapping from the fibre over $\alpha(0)$ into the one over $\alpha(1)$ satisfying some axioms. [22, sec. 2.2]
1964	C. Teleman	Definition of a connection in topological bundle as a parallel transport along canonical paths $\alpha: [0, 1] \rightarrow M$ in its base M . The parallel transport is defined as a lifting of these paths through a point in the fibre over their initial points $\alpha(0)$. [24, chapter IV, sec. B.3]
1968	P. Dombrowski	Definition of a linear connection in vector bundle as a parallel transport along paths $\beta: [a, b] \rightarrow M$, with $a, b \in \mathbb{R}$ and $a \leq b$, in its base M . The parallel transport is defined as a mapping from the fibre over $\alpha(a)$ into the one over $\alpha(b)$ satisfying certain axioms. The theory of covariant derivatives is constructed on that base. [25, § 1]
1981	Walter Poor	A detailed axiomatic definition of a parallel transport in vector bundles. The whole theory of linear connections in such bundles is deduced on that ground. [32]

The purpose of the present investigation is to be revealed different relations between the concepts “connection on a bundle” and “(parallel) transport (along paths) in a bundle”. These concepts will be shown to be equivalent in a sense that there exist bijective mappings between the sets of connections, (axiomatically defined) parallel transports and transports along paths satisfying some additional conditions. In this way, two different, but equivalent, systems of axioms defining the concept “parallel transport” will be established.

The paper is organized as follows.

Section 2 recalls the definition of a connection on a bundle as a distribution on its bundle space which is complimentary to the vertical distribution on it. The notions of parallel transport generated by connection and of (axiomatically defined) parallel transport are introduced. In section 3 is defined the notion of a transport along paths, which is one of the possible axiomatic approaches to the concept of a “parallel transport”. Section 4 shows how a transport along paths generates a connection and presents some properties of such a connection. The main result of section 5 is the construction of bijective mapping between the sets of transports along paths, satisfying some additional conditions, and parallel transports. Section 6 deals with different relations between connections, parallel transports, and transports along paths. The equivalence between the concepts connection, transport along

paths, satisfying some additional conditions, and parallel transport is established. Section 7 summarizes the basic results of the present work in a series of theorems. In section 8 some of the preceeding results are generalized in a case of general topological bundles. In section 9 are proved some results which are specific for vector bundles. In particular, a conditions ensuring equivalence between linear transport along paths and a linear connection in these bundles is presented. Section 10 closes the paper with some concluding remarks.

Let us now present some introductory material, like notation etc., that will be needed for our exposition. The reader is referred for details to standard books on differential geometry, like [5, 32, 34].

A differentiable finite-dimensional manifold over a field $\mathbb{K}$ will be denoted typically by M . Here $\mathbb{K}$ stands for the field $\mathbb{R}$ of real or the field $\mathbb{C}$ of complex numbers, $\mathbb{K} = \mathbb{R}, \mathbb{C}$. The manifolds we consider are supposed to be smooth of class C^1 .^a

The set of vector fields, realized as first order differential operators, over M will be denoted by $\mathcal{X}(M)$. The space tangent to M at $p \in M$ is $T_p(M)$ and $(T(M), \pi_T, M)$ stand for the tangent bundle over M . The value of $X \in \mathcal{X}(M)$ at $p \in M$ is $X_p \in T_p(M)$.

By $J \subseteq \mathbb{R}$ will be denoted an arbitrary real interval that can be open or closed at one or both its ends. The notation $[\sigma, \tau]$ will be used for an arbitrary closed interval with ends $\sigma, \tau \in \mathbb{R}$, with $\sigma \leq \tau$. The notation $\gamma: J \rightarrow M$ represents an arbitrary path in M . For a C^1 path $\gamma: J \rightarrow M$, the vector tangent to γ at $s \in J$ will be denoted by $\dot{\gamma}(s) := \left. \frac{d}{dt} \right|_{t=s} (\gamma(t)) \in T_{\gamma(s)}(M)$. If $s_0 \in J$ is an end point of J and J is closed at s_0 , the derivative in the definition of $\dot{\gamma}(s_0)$ is regarded as a one-sided derivative at s_0 .

In this work, we shall need the notion of an inverse path and of a product of paths. There are not ‘natural’ definitions of these concepts, but this is not important for us as the (parallel) transports we shall consider bellow are parametrization invariant in some sense, like (3.5) below. For that reason, the concepts mentioned will be defined only for *canonical paths* $[0, 1] \rightarrow M$, whose domain is the real interval $[0, 1] := \{r \in \mathbb{R} : 0 \leq r \leq 1\}$. The path *inverse* to $\gamma: [0, 1] \rightarrow M$ is $\gamma_- := \gamma \circ \tau_-: [0, 1] \rightarrow M$, with $\circ$ being the sign of composition of mappings and $\tau_-: [0, 1] \rightarrow [0, 1]$ being given by $\tau_-(t) := 1 - t$ for $t \in [0, 1]$. If $\gamma_1, \gamma_2: [0, 1] \rightarrow M$ and $\gamma_1(1) = \gamma_2(0)$, the *product* $\gamma_1\gamma_2$ of γ_1 and γ_2 is a canonical path $\gamma_1\gamma_2: [0, 1] \rightarrow M$ such that $(\gamma_1\gamma_2)(t) := \gamma_1(2t)$ for $t \in [0, 1/2]$ and $(\gamma_1\gamma_2)(t) := \gamma_2(2t - 1)$ for $t \in [1/2, 1]$. For more details on this item, see [35, 36].

If $k \in \mathbb{N}$ and $k \leq \dim M$, a k -dimensional *distribution* Δ on M is defined as a mapping $\Delta: p \mapsto \Delta_p$ assigning to each $p \in M$ a k -dimensional subspace Δ_p of the tangent space $T_p(M)$ of M at p , $\Delta_p \subseteq T_p(M)$. We say that a vector field $X \in \mathcal{X}(M)$ is in Δ and write $X \in \Delta$, if $X_p \in \Delta_p$ for all $p \in M$.

By (E, π, M) , we shall denote a bundle with bundle space E , projection $\pi: E \rightarrow M$, and base space M . Suppose that the spaces M and E are manifolds of finite dimensions $n \in \mathbb{N}$ and $n + r$, for some $r \in \mathbb{N}$, respectively; so the dimension of the fibre $\pi^{-1}(x)$, with $x \in M$, i.e. the fibre dimension of (E, π, M) , is r . Besides, let these manifolds be C^1 differentiable, if the opposite is not stated explicitly.^b

^a Some of our definitions or/and results are valid also for C^0 manifolds, but we do not want to overload the material with continuous counting of the required degree of differentiability of the manifolds involved. Some parts of the text admit generalizations on more general spaces, like the topological ones, but this is out of the subject of the present work.

^b Some of our considerations are valid also for C^0 manifolds. By assuming C^1 differentiability, we skip the problem of counting the required differentiability class of the whole material that follows. Sometimes, the C^1 differentiability is required explicitly, which is a hint that a statement or definition is not valid otherwise.

2. Connections and the generated by them parallel transports (review)

From a number of equivalent definitions of a connection on differentiable manifold [21, sections 2.1 and 2.2], we shall use the following one.

Definition 2.1. A *connection on a bundle* (E, π, M) is an $n = \dim M$ dimensional distribution Δ^h on E such that, for each $p \in E$ and the *vertical distribution* Δ^v defined by

$$\Delta^v: p \mapsto \Delta_p^v := T_{\iota(p)}(\pi^{-1}(\pi(p))) \cong T_p(\pi^{-1}(\pi(p))), \quad (2.1)$$

with $\iota: \pi^{-1}(\pi(p)) \rightarrow E$ being the inclusion mapping, is fulfilled

$$\Delta_p^v \oplus \Delta_p^h = T_p(E), \quad (2.2)$$

where $\Delta^h: p \mapsto \Delta_p^h \subseteq T_p(E)$ and $\oplus$ is the direct sum sign. The distribution Δ^h is called *horizontal* and symbolically we write $\Delta^v \oplus \Delta^h = T(E)$.

A *vector* at a point $p \in E$ (resp. a *vector field* on E) is said to be *vertical* or *horizontal* if it (resp. its value at p) belongs to Δ_p^v or Δ_p^h , respectively, for the given (resp. any) point p . A vector $Y_p \in T_p(E)$ (resp. vector field $Y \in \mathcal{X}(E)$) is called a *horizontal lift of a vector* $X_{\pi(p)} \in T_{\pi(p)}(M)$ (resp. *vector field* $X \in \mathcal{X}(M)$ on $M = \pi(E)$) if $\pi_*(Y_p) = X_{\pi(p)}$ for the given (resp. any) point $p \in E$. Since $\pi_*|_{\Delta_p^h}: \Delta_p^h \rightarrow T_{\pi(p)}(M)$ is a vector space isomorphism for all $p \in E$ [32, sec. 1.24], any vector in $T_{\pi(p)}(M)$ (resp. vector field in $\mathcal{X}(M)$) has a unique horizontal lift in $T_p(E)$ (resp. $\mathcal{X}(E)$).

As a result of (2.2), any vector $Y_p \in T_p(E)$ (resp. vector field $Y \in \mathcal{X}(E)$) admits a unique representation $Y_p = Y_p^v \oplus Y_p^h$ (resp. $Y = Y^v \oplus Y^h$) with $Y_p^v \in \Delta_p^v$ and $Y_p^h \in \Delta_p^h$ (resp. $Y^v \in \Delta^v$ and $Y^h \in \Delta^h$). If the distribution $p \mapsto \Delta_p^h$ is differentiable of class C^m , $m \in \mathbb{N} \cup \{0\}$, it is said that the *connection* Δ^h is (differentiable) of class C^m . A connection Δ^h is of class C^m if and only if, for every C^m vector field Y on E , the vertical Y^v and horizontal Y^h vector fields are of class C^m .

A C^1 path $\beta: J \rightarrow E$ is called *horizontal* (*vertical*) if its tangent vector $\dot{\beta}$ is horizontal (vertical) vector along β , i.e. $\dot{\beta}(s) \in \Delta_{\beta(s)}^h$ ($\dot{\beta}(s) \in \Delta_{\beta(s)}^v$) for all $s \in J$. A lift $\bar{\gamma}: J \rightarrow E$ of a path $\gamma: J \rightarrow M$, i.e. $\pi \circ \bar{\gamma} = \gamma$, is called *horizontal* if $\bar{\gamma}$ is a horizontal path, i.e. when the vector field $\dot{\bar{\gamma}}$ tangent to $\bar{\gamma}$ is horizontal or, equivalently, if $\dot{\bar{\gamma}}$ is a horizontal lift of $\dot{\gamma}$. Since $\pi^{-1}(\gamma(J))$ is an $(r+1)$ dimensional submanifold of E , the distribution $p \mapsto \Delta_p^h \cap T_p(\pi^{-1}(\gamma(J)))$ is one-dimensional and, consequently, is integrable. The integral paths of that distribution are horizontal lifts of γ and, for each $p \in \pi^{-1}(\gamma(J))$, there is a unique horizontal lift $\bar{\gamma}_p$ of γ passing through p .

Definition 2.2. Let $\gamma: [\sigma, \tau] \rightarrow M$, with $\sigma, \tau \in \mathbb{R}$ and $\sigma \leq \tau$, and $\bar{\gamma}_p$ be the unique horizontal lift of γ in E passing through $p \in \pi^{-1}(\gamma([\sigma, \tau]))$. The *parallel transport (translation, displacement)* generated by (assigned to, defined by) a connection Δ^h is a mapping $P: \gamma \mapsto P^\gamma$, assigning a mapping to the path γ

$$P^\gamma: \pi^{-1}(\gamma(\sigma)) \rightarrow \pi^{-1}(\gamma(\tau)) \quad \gamma: [\sigma, \tau] \rightarrow M \quad (2.3)$$

such that, for each $p \in \pi^{-1}(\gamma(\sigma))$,

$$P^\gamma(p) := \bar{\gamma}_p(\tau). \quad (2.4)$$

Recall now the basic properties of the parallel transports generated by connections.

Proposition 2.1. *Let*

$$P: \gamma \mapsto P^\gamma: \pi^{-1}(\gamma(\sigma)) \rightarrow \pi^{-1}(\gamma(\tau)) \quad \gamma: [\sigma, \tau] \rightarrow M \quad (2.5)$$

be the parallel transport generated by a connection on some bundle (E, π, M) . The mapping P has the following properties:

(i) The parallel transport P is invariant under orientation preserving changes of the paths' parameters. Precisely, if $\gamma: [\sigma, \tau] \rightarrow M$ and $\chi: [\sigma', \tau'] \rightarrow [\sigma, \tau]$ is an orientation preserving C^1 diffeomorphism, then

$$P^{\gamma \circ \chi} = P^\gamma. \quad (2.6)$$

(ii) If $\gamma: [0, 1] \rightarrow M$ and $\gamma_-: [0, 1] \rightarrow M$ is its canonical inverse, $\gamma_-(t) = \gamma(1 - t)$ for $t \in [0, 1]$, then

$$P^{\gamma_-} = (P^\gamma)^{-1}. \quad (2.7)$$

(iii) If $\gamma_1, \gamma_2: [0, 1] \rightarrow M$, $\gamma_1(1) = \gamma_2(0)$, and $\gamma_1\gamma_2: [0, 1] \rightarrow M$ is their canonical product, then

$$P^{\gamma_1\gamma_2} = P^{\gamma_2} \circ P^{\gamma_1}. \quad (2.8)$$

(iv) If $\gamma_{r,x}: \{r\} = [r, r] \rightarrow \{x\}$ for some given $r \in \mathbb{R}$ and $x \in M$, then

$$P^{\gamma_{r,x}} = \text{id}_{\pi^{-1}(x)}. \quad (2.9)$$

(v) If $\bar{\gamma}_p: s \in [\sigma, \tau] \rightarrow \bar{\gamma}(s) := P^{\gamma|[\sigma, s]}(p)$ is the lifting of γ through $p \in \pi^{-1}(\gamma(\sigma))$ defined by P and the C^1 paths $\gamma_i: [\sigma_i, \tau_i] \rightarrow M$ are such that $\gamma_1(\sigma_1) = \gamma_2(\sigma_2)$ and $\dot{\gamma}_1(\sigma_1) = \dot{\gamma}_2(\sigma_2)$, then (a) the lifted path

$$\bar{\gamma}_p \text{ is of class } C^1 \quad (2.10)$$

for any C^1 path γ and (b) the lifted paths $\bar{\gamma}_{1;p}$ and $\bar{\gamma}_{2;p}$ have equal tangent vectors at p ,

$$\dot{\bar{\gamma}}_{1;p}(\sigma_1) = \dot{\bar{\gamma}}_{2;p}(\sigma_2). \quad (2.11)$$

(vi) If $\gamma_i: [\sigma_i, \tau_i] \rightarrow M$, $i = 1, 2$, are two C^1 paths and $\gamma_1(\sigma_1) = \gamma_2(\sigma_2)$, then for every $a_1, a_2 \in \mathbb{K}$ there exists a C^1 path $\gamma_3: [\sigma_3, \tau_3] \rightarrow M$ such that $\gamma_3(\sigma_3) = \gamma_1(\sigma_1) (= \gamma_2(\sigma_2))$ and the vector tangent to the lifted path $\bar{\gamma}_{3;p}: t_3 \in [\sigma_3, \tau_3] \rightarrow \bar{\gamma}_{3;p}(t_3) := P^{\gamma_3|[\sigma_3, t_3]}(p)$, with $p \in \pi^{-1}(\gamma_3(\sigma_3))$, at σ_3 is

$$\dot{\bar{\gamma}}_{3;p}(\sigma_3) = a_1 \dot{\bar{\gamma}}_{1;p}(\sigma_2) + a_2 \dot{\bar{\gamma}}_{2;p}(\sigma_2), \quad (2.12)$$

where $\bar{\gamma}_{i;p}: t_i \in [\sigma_i, \tau_i] \rightarrow \bar{\gamma}_{i;p}(t_i) := P^{\gamma_i|[\sigma_i, t_i]}(p)$.

Remark 2.1. As a result of (2.6), some properties of the parallel transports generated by connections, like (2.7) and (2.8), are sufficient to be formulated/proved only for canonical paths $[0, 1] \rightarrow M$.

Proof. The proofs of (2.6)–(2.9) can be found in a number of works, for example in [2, 4, 22, 26, 28, 29, 31, 37–39] $\square$

Definition 2.3. A mapping (2.5) satisfying (2.6)–(2.12) will be called (axiomatically defined) parallel transport.

Proposition 2.2. Let P be the parallel transport assigned to a C^m , with $m \in \mathbb{N} \cup \{0\}$, connection on a smooth, of class C^{m+1} , bundle (E, π, M) . Then P is smooth, of class C^m , in a sense that, if $\gamma: [\sigma, \tau] \rightarrow M$ is a C^1 path, then P^γ is in the set of C^m diffeomorphisms between the fibres $\pi^{-1}(\gamma(\sigma))$ and $\pi^{-1}(\gamma(\tau))$,

$$P: \gamma \mapsto P^\gamma \in \text{Diff}^m(\pi^{-1}(\gamma(\sigma)), \pi^{-1}(\gamma(\tau))) \quad \gamma: [\sigma, \tau] \rightarrow M. \quad (2.13)$$

Proof. See [8, 11, 14]. □

The axiomatic approach to parallel transport was developed mainly on the ground on the properties (2.6)–(2.13) of the parallel transports assigned to connections. However, this topic is out of the range of the present paper and the reader is referred to the literature cited at the beginning of section 3.

3. Transports along paths in fibre bundles (review)

The widespread approach to the concept of a “parallel transport” is it to be considered as a secondary one and defined on the basis of the connection theory [2–19]. However, the opposite approach, in which the parallel transport is axiomatically defined and from it the connection theory is constructed, is also known [22–32] and goes back to 1949^c; e.g. it is systematically realized in [32], where the connection theory on vector bundles is investigated. In [41] the concept of a “parallel transport” was generalize to the one of “transport along paths”. The relations between both concepts were analyzed in [42]; in particular, theorem 3.1 of [42, p. 13] (see theorem 5.1 below) contains a necessary and sufficient condition for a transport along paths to be (axiomatically defined) parallel transport.

Definition 3.1. A *transport along paths* in a (topological) bundle (E, π, B) is a mapping I assigning to every path $\gamma: J \rightarrow M$ a mapping I^γ , termed *transport along γ* , such that $I^\gamma: (s, t) \mapsto I_{s \rightarrow t}^\gamma$ where the mapping

$$I_{s \rightarrow t}^\gamma: \pi^{-1}(\gamma(s)) \rightarrow \pi^{-1}(\gamma(t)) \quad s, t \in J, \quad (3.1)$$

called *transport along γ from s to t* , has the properties:

$$I_{s \rightarrow t}^\gamma \circ I_{r \rightarrow s}^\gamma = I_{r \rightarrow t}^\gamma \quad r, s, t \in J \quad (3.2)$$

$$I_{s \rightarrow s}^\gamma = \text{id}_{\pi^{-1}(\gamma(s))} \quad s \in J, \quad (3.3)$$

where $\circ$ denotes composition of mappings and id_X is the identity mapping of a set X .

An analysis and various comments on this definition can be found in [41–44].

As we shall see below, an important role is played by transports along paths satisfying some additional conditions, in particular

$$I_{s \rightarrow t}^{\gamma|J'} = I_{s \rightarrow t}^\gamma \quad s, t \in J' \quad (3.4)$$

$$I_{s \rightarrow t}^{\gamma \circ \chi} = I_{\chi(s) \rightarrow \chi(t)}^\gamma \quad s, t \in J'', \quad (3.5)$$

where $J' \subseteq J$ is a subinterval, $\gamma|J'$ is the restriction of γ to J' , and $\chi: J'' \rightarrow J$ is a bijection of a real interval J'' onto J .

Putting $r = t$ in (3.2) and using (3.3), we see that the mappings (3.1) are invertible and

$$(I_{s \rightarrow t}^\gamma)^{-1} = I_{t \rightarrow s}^\gamma. \quad (3.6)$$

For discussion and more details on the transports along paths, the reader is referred to [41, 42]; in particular, there can be found the general form of these mappings, possible restrictions on them, their relations with other differential-geometric structures, etc.

^c It seems that the earliest written accounts on this approach are the ones due to Ü. G. Lumiste [22, sec. 2.2] and C. Teleman [24, chapter IV, sec. B.3] (both published in 1964), the next essential steps being made by P. Dombrowski [25, § 1] and W. Poor [32]. Besides, the author of [25] states that his paper is based on unpublished lectures of prof. Willi Rinow (1907–1979) in 1949; see also [32, p. 46] where the author claims that the first axiomatical definition of a parallel transport in the tangent bundle case is given by prof. W. Rinow in his lectures at the Humboldt University in 1949. Some heuristic comments on the axiomatic approach to parallel transport theory can be found in [40, sec. 2.1] too.

4. A connection generated by transport along paths

The following result describes how a transport along paths generates a connection.^d

Theorem 4.1. *Let I be a transport along paths in a bundle (E, π, M) . Let $\gamma: J \rightarrow M$ be a path and, for any $s_0 \in J$ and $p \in \pi^{-1}(\gamma(s_0))$, the lift $\bar{\gamma}_{s_0,p}: J \rightarrow E$ of γ be defined by*

$$\bar{\gamma}_{s_0,p}(t) = I_{s_0 \rightarrow t}^\gamma(p) \quad t \in J. \quad (4.1)$$

Suppose the transport I is such that:

(a) (C^1 smoothness) The path

$$\bar{\gamma}_{s_0,p} \text{ is of class } C^1 \quad (4.2a)$$

for every C^1 path γ , s_0 and p .

(b) (Initial uniqueness) If $\gamma_i: J_i \rightarrow M$, $i = 1, 2$, are two C^1 paths and for some $s_i \in J_i$ is fulfilled $\gamma_1(s_1) = \gamma_2(s_2)$ and $\dot{\gamma}_1(s_1) = \dot{\gamma}_2(s_2)$, then the lifted paths $\bar{\gamma}_{i;s_i,p}$, defined via (4.1) with $p \in \pi^{-1}(\gamma_1(s_1)) = \pi^{-1}(\gamma_2(s_2))$, have equal tangent vectors at p ,

$$\dot{\bar{\gamma}}_{1;s_1,p}(s_1) = \dot{\bar{\gamma}}_{2;s_2,p}(s_2). \quad (4.2b)$$

(c) (Linearization) If $\gamma_i: J_i \rightarrow M$, $i = 1, 2$, are two C^1 paths and for some $s_i \in J_i$ is fulfilled $\gamma_1(s_1) = \gamma_2(s_2)$, then for every $a_1, a_2 \in \mathbb{K}$ there exists a C^1 path $\gamma_3: J_3 \rightarrow M$ (generally depending on a_1, a_2, γ_1 and γ_2) such that $\gamma_3(s_3) = \gamma_1(s_1) (= \gamma_2(s_2))$ for some $s_3 \in J_3$ and the vector tangent to the lifted path $\bar{\gamma}_{3;s_3,p}$, defined via (4.1) with $p \in \pi^{-1}(\gamma_3(s_3))$, at s_3 is

$$\dot{\bar{\gamma}}_{3;s_3,p}(s_3) = a_1 \dot{\bar{\gamma}}_{1;s_1,p}(s_1) + a_2 \dot{\bar{\gamma}}_{2;s_2,p}(s_2). \quad (4.2c)$$

Then

$$\begin{aligned} \Delta^I: p \mapsto \Delta_p^I &:= \left\{ \frac{d}{dt} \Big|_{t=s_0} (\bar{\gamma}_{s_0,p}(t)) \right. \\ &\quad \left. : \gamma: J \rightarrow M \text{ is } C^1 \text{ and injective, } s_0 \in J, \gamma(s_0) = \pi(p) \right\} \subseteq T_p(E), \end{aligned} \quad (4.3)$$

with $p \in E$, is a distribution which is a connection on (E, π, M) , i.e.

$$\Delta_p^v \oplus \Delta_p^I = T_p(E) \quad p \in E, \quad (4.4)$$

with Δ^v being the vertical distribution on E , $\Delta_p^v = T_p(\pi^{-1}(\pi(p)))$.

Proof. To begin with, we shall write the following result.

Lemma 4.1. *If a C^1 path $\bar{\gamma}: J \rightarrow E$ is a lift of a C^1 path $\gamma: J \rightarrow M$, $\pi \circ \bar{\gamma} = \gamma$, then*

$$\pi_*(\dot{\bar{\gamma}}(t)) = \dot{\gamma}(t). \quad (4.5)$$

Proof of Lemma 4.1 Let $\{u^\mu = x^\mu \circ \pi, u^a\}$ be bundle coordinate system on E and p be a point in its domain. Then (4.5) follows from $\pi_*\left(\frac{\partial}{\partial u^I} \Big|_p\right) = \frac{\partial(x^\mu \circ \pi)}{\partial u^I} \Big|_p \frac{\partial}{\partial x^\mu} \Big|_{\pi(p)}$ and

$$\dot{\bar{\gamma}}^\mu = \dot{\gamma}^\mu \quad (4.6)$$

which is a corollary of $\dot{\bar{\gamma}}^\mu(t) = \frac{d(u^\mu \circ \bar{\gamma}(t))}{dt} = \frac{d(x^\mu \circ \gamma(t))}{dt} = \dot{\gamma}^\mu(t)$ for all $t \in J$. $\square$

^d The author thanks Petko Nikolov (Physical department of the Sofia University “St. Kliment Ohridski”) for a discussion which led to a correct formulation of theorem 4.1.

From (4.3) and (4.5), we get

$$\pi_*(\Delta_p^I) = \{\dot{\gamma}(s_0) : \gamma : J \rightarrow M \text{ is } C^1 \text{ and injective, } s_0 \in J, \gamma(s_0) = \pi(p)\} = T_{\pi(p)}(M)$$

as $\dot{\gamma}(s_0)$ is an arbitrary vector in $T_{\pi(p)}(M) = T_{\gamma(s_0)}(M)$. Besides, the condition (c) ensures that Δ_p^I is a vector subspace in $T_p(E)$. Thus $\pi_*|_{\Delta_p^I} : \Delta_p^I \rightarrow T_{\pi(p)}(M)$ is a surjective mapping between vector spaces. It is also linear as it is a restriction of the tangent mapping π_* , which in turn is a linear mapping [5, sec. 1.22], on the vector subspace Δ_p^I . At last, we shall prove that $\pi_*|_{\Delta_p^I}$ is injective, from where it follows that $\pi_*|_{\Delta_p^I}$ is a vector space isomorphism for every $p \in E$ which, in its turn, implies that $\Delta^I : p \rightarrow \Delta_p^I$ is a distribution satisfying (4.4) as $\pi_*(\Delta_p^v) = 0_{\pi(p)} \in T_{\pi(p)}(M)$.

If $G_i \in \Delta_p^I$, $i = 1, 2$, then there exist paths $\gamma_i : J_i \rightarrow M$ such that $\gamma_i(s_i) = \pi(p)$ for some $s_i \in J_i$ and $G_i = \dot{\gamma}_{i;s_i,p}(s_i)$, with $i = 1, 2$ and the lifted paths in the r.h.s. being given by (4.1). Then $\pi_*(G_i) = \dot{\gamma}_i(s_i)$, due to (4.5). Suppose that $\dot{\gamma}_1(s_1) = \dot{\gamma}_2(s_2)$. Then condition (b) entails $\dot{\gamma}_{1;s_1,p}(s_1) = \dot{\gamma}_{2;s_2,p}(s_2)$ so that $G_1 = G_2$, which means that $\pi_*|_{\Delta_p^I} : \Delta_p^I \rightarrow T_{\pi(p)}(M)$ is injective. $\square$

Remark 4.1. The condition (a) ensures that all our constructions have a sense. If the condition (b) is not valid, then $\pi_*|_{\Delta_p^I}$ is linear and surjective, but we cannot prove that it is also injective; precisely, $\pi_*|_{\Delta_p^h}$ is injective iff the property (b) is valid. At last, the condition (c) guarantees that Δ_p^I is a vector subspace of $T_p(E)$; in fact, Δ_p^I is a vector space iff the property (c) is valid. This is quite an essential moment as the direct complement B to a vector subspace A of a vector space V (in our case $B = \Delta_p^h$, $A = \Delta_p^v$ and $V = T_p(E)$) is generally *not* a vector subspace; i.e. the equality $V = A \oplus B$, with V and A being vector spaces, does not generally imply that B is a vector space; for example, if $A = \{(x, 0) : x \in \mathbb{R}\} \subset \mathbb{R}^2$, then $\mathbb{R}^2 = A \oplus B$ with $B = \{(x, y) \in \mathbb{R}^2, y \neq 0\} \cup \{(0, 0)\}$, but B is not a vector subspace in $\mathbb{R}^2$, as $(a, -c), (b, +c) \in B$ for $a, b, c \in \mathbb{R}$ and $c \neq 0$ but $(a, -c) + (b, +c) = (a + b, 0) \notin B$ for $a + b \neq 0$.

Remark 4.2. Since (4.2c) and (4.5) entail

$$\dot{\gamma}_3(s_3) = a_1 \dot{\gamma}_1(s_1) + a_2 \dot{\gamma}_2(s_2), \quad (4.7a)$$

the path γ_3 is uniquely fixed by the conditions that it passes through the point $\pi(p)$,

$$\gamma_3(s_3) = \pi(p). \quad (4.7b)$$

Therefore, condition (c) is equivalent to the requirement the vector tangent at s_3 to the lift, given via (4.1), of the path defined via (4.7) to be (4.2c).

Remark 4.3. The role of the transport I along paths in theorem 4.1 is on its base to be constructed a lifting of the paths in M to paths in E with appropriate properties. Namely, such a lifting should assign to a path $\gamma : J \rightarrow M$ a unique path $\bar{\gamma}_{s_0,p} : J \rightarrow E$ passing through a given point $p \in \pi^{-1}(\gamma(s_0))$, for some $s_0 \in J$, and such that $\pi \circ \bar{\gamma}_{s_0,p} = \gamma$ and, if $q \in \bar{\gamma}_{s_0,p}(t_0)$ for some $t_0 \in J$, then $\bar{\gamma}_{t_0,q} = \bar{\gamma}_{s_0,p}$. On this ground one can generalize theorem 4.1 as well as some of the next considerations and results.

Definition 4.1. The connection Δ^I , defined in theorem 4.1, will be called *assigned to (defined by, generated by)* the transport I along paths.

Remark 4.4. Note that the transport I in this definition must have the properties (4.2) described in theorem 4.1.

It is worth recording the simple fact that, if equation (3.4) or (3.5) holds, then the path $\bar{\gamma}_{s_0,p}$, defined via (4.1) is such that respectively

$$\bar{\gamma}_{s_0,p}|_{J'} = (\overline{\gamma|J'})_{s_0,p} \quad s_0 \in J' \quad (4.8)$$

$$\bar{\gamma}_{\chi(s_0),p} \circ \chi = (\overline{\gamma \circ \chi})_{s_0,p} \quad s_0 \in J''. \quad (4.9)$$

Proposition 4.1. *For C^1 paths, the equations (3.4) and (3.5) are consequences of the suppositions (a)–(c) from theorem 4.1 but the converse is generally not true, i.e. (3.4) and (3.5) are necessary but generally not sufficient conditions for (4.3) to be a connection.*

Proof. To prove (3.4), use assumption (b) and the local existence of a unique C^1 path passing through a given point and having a fixed tangent vector at it. The proof of (3.5) is more complicated. For the purpose, observe that the paths $\bar{\beta}_{s_0,p}: t \rightarrow I_{s_0 \rightarrow t}^\beta(p)$ and $\bar{\delta}_{s_0,p}: t \rightarrow I_{\chi(s_0) \rightarrow \chi(t)}^\gamma(p) = \bar{\gamma}_{\chi(s_0),p} \circ \chi(t)$, for $s_0, t \in J''$, in E are liftings of the path $\beta = \gamma \circ \chi$ and are Δ^I -horizontal, i.e. $\dot{\bar{\beta}}_{s_0,p}(t) \in \Delta_{\bar{\beta}_{s_0,p}(t)}^I$ and $\dot{\bar{\delta}}_{s_0,p}(t) \in \Delta_{\bar{\delta}_{s_0,p}(t)}^I$. (Here use that $\dot{\beta}(t) = \frac{d\chi(t)}{dt} \dot{\gamma}(\chi(t))$ and similarly for $\bar{\delta}_{s_0,p}$.) Then the existence of a unique horizontal lift of any C^1 path (see also proposition 4.2 below) implies $\bar{\beta}_{s_0,p} = \bar{\delta}_{s_0,p}$. $\square$

Proposition 4.2. *If $\gamma: J \rightarrow M$ is an injective path and $p \in \pi^{-1}(\gamma(s_0))$ for some $s_0 \in J$, then there is a unique Δ^I -horizontal lift of γ (relative to Δ^I) in E through p and it is exactly the path $\bar{\gamma}_{s_0,p}$ defined by (4.1).*

Proof. Simply apply the definitions (4.1) and (4.3) and use the properties (3.1) and (3.2) of the transports along paths ($t_0 \in J$):

$$\begin{aligned} \dot{\bar{\gamma}}_{s_0,p}(t_0) &= \frac{d}{dt} \Big|_{t=t_0} (I_{s_0 \rightarrow t}^\gamma(p)) = \frac{d}{dt} \Big|_{t=t_0} (I_{t_0 \rightarrow t}^\gamma(I_{s_0 \rightarrow t_0}^\gamma(p))) \\ &= \frac{d}{dt} \Big|_{t=t_0} (I_{t_0 \rightarrow t}^\gamma(\bar{\gamma}_{s_0,p}(t_0))) = \dot{\bar{\gamma}}_{t_0, \bar{\gamma}_{s_0,p}(t_0)}(t_0) \in \Delta_{\bar{\gamma}_{s_0,p}(t_0)}^I. \end{aligned}$$

$\square$

Remark 4.5. If γ is not injective and $\gamma(s_0) = \gamma(t_0)$ for some $s_0, t_0 \in J$ such that $s_0 \neq t_0$, then the paths $\bar{\gamma}_{s_0,p}, \bar{\gamma}_{t_0,p}: J \rightarrow E$ need not to coincide as $\bar{\gamma}_{s_0,p} = I_{t_0 \rightarrow s_0} \circ \bar{\gamma}_{t_0,p}$, due to (4.1) and (3.2). Therefore the Δ^I -horizontal lift of a non-injective path through a point in E lying in a fibre over a self-intersection point of the path, if any, may not be unique. Similar is the situation for an arbitrary connection. (This range of problems is connected with the so-called holonomy groups.)

Proposition 4.3. *The parallel transport l generated by the connection, defined by a transport along paths I , is such that*

$$\mathsf{l}^\gamma = I_{\sigma \rightarrow \tau}^\gamma \quad \text{for } \gamma: [\sigma, \tau] \rightarrow M. \quad (4.10)$$

Proof. According to definition 2.2 and (4.1), we have:

$$\begin{aligned} \mathsf{l}: \gamma &\mapsto \mathsf{l}^\gamma: \pi^{-1}(\gamma(\sigma)) \rightarrow \pi^{-1}(\gamma(\tau)) & \gamma: [\sigma, \tau] &\rightarrow M \\ \mathsf{l}: p &\mapsto \mathsf{l}^\gamma(p) = \bar{\gamma}_{\sigma,p}(\tau) = I_{\sigma \rightarrow \tau}^\gamma(p) & p &\in \pi^{-1}(\gamma(\sigma)). \end{aligned}$$

$\square$

5. Links between parallel transports and transports along paths

Now we shall present a modified version of [42, p. 13, theorem 3.1].

Theorem 5.1. *Let I be a transport along paths in a bundle (E, π, M) and $\gamma: [\sigma, \tau] \rightarrow M$. If I satisfies the conditions (4.2) from theorem 4.1, then the mapping*

$$\mathsf{l}: \gamma \mapsto \mathsf{l}^\gamma := I_{\sigma \rightarrow \tau}^\gamma: \pi^{-1}(\gamma(\sigma)) \rightarrow \pi^{-1}(\gamma(\tau)) \quad \gamma: [\sigma, \tau] \rightarrow M \quad (5.1)$$

is a parallel transport, i.e. it possess the properties (2.6)–(2.12), with $\mathbf{l}$ for $\mathbf{P}$. Besides, if I is smooth in a sense that

$$I_{s \rightarrow t}^\beta \in \text{Diff}^m(\pi^{-1}(\beta(s)), \pi^{-1}(\beta(t))) \quad \beta: J \rightarrow M \quad s, t \in J \quad (5.2)$$

for some $m \in \mathbb{N} \cup \{0\}$, then the mapping (5.1) satisfies (2.13), with $\mathbf{l}$ for $\mathbf{P}$.

Conversely, suppose the mapping

$$\mathbf{P}: \gamma \mapsto \mathbf{P}^\gamma: \pi^{-1}(\gamma(\sigma)) \rightarrow \pi^{-1}(\gamma(\tau)) \quad \gamma: [\sigma, \tau] \rightarrow M \quad (5.3)$$

is a parallel transport, i.e. satisfies (2.6)–(2.12), and define the mapping

$$P: \beta \mapsto P^\beta: (s, t) \mapsto P_{s \rightarrow t}^\beta = \mathbf{P}^{(\beta|[\sigma, \tau]) \circ \chi_t^{[\sigma, \tau]}} \circ \left(\mathbf{P}^{(\beta|[\sigma, \tau]) \circ \chi_s^{[\sigma, \tau]}} \right)^{-1} \quad \beta: J \rightarrow M, \quad (5.4)$$

where $s, t \in J$, $\sigma, \tau \in J$ are such that $\sigma \leq \tau$ and $[\sigma, \tau] \ni s, t$,^e and $\chi_s^{[\sigma, \tau]}: [\sigma, \tau] \rightarrow [\sigma, s]$ are for $s > \sigma$ arbitrary orientation preserving C^1 diffeomorphisms (depending on β via the interval $[\sigma, \tau]$). Then the mapping (5.4) is a transport along paths in (E, π, M) , which transport satisfies the conditions (4.2), with P for I . Besides, under the same assumptions, the condition (2.13) for $\mathbf{P}$ implies (5.2), with P for I , where P is given by (5.4).

Remark 5.1. Instead by (5.4), the transport P along paths generated by a parallel transport $\mathbf{P}$ can be defined equivalently as follows. For a path $\gamma: [\sigma, \tau] \rightarrow M$ and $s, t \in [\sigma, \tau]$, we put (see (2.6))

$$P: \gamma \mapsto P^\gamma: (s, t) \mapsto P_{s \rightarrow t}^\gamma = \mathbf{P}^{\gamma \circ \chi_t^{[\sigma, \tau]}} \circ \left(\mathbf{P}^{\gamma \circ \chi_s^{[\sigma, \tau]}} \right)^{-1} = \mathbf{P}^{\gamma|[\sigma, t]} \circ \left(\mathbf{P}^{\gamma|[\sigma, s]} \right)^{-1} \quad \gamma: [\sigma, \tau] \rightarrow M \quad (5.5a)$$

Now, for an arbitrary path $\beta: J \rightarrow M$, with J being closed or open at one or both its ends, we set

$$P: \beta \mapsto P^\beta: (s, t) \mapsto P_{s \rightarrow t}^\beta = \begin{cases} P_{s \rightarrow t}^{\beta|[\sigma, t]} & \text{for } s \leq t \\ (P_{t \rightarrow s}^{\beta|[\sigma, s]})^{-1} & \text{for } s \geq t \end{cases} \quad \beta: J \rightarrow M \quad s, t \in J. \quad (5.5b)$$

It can easily be verified that (5.5) are tantamount to

$$P: \beta \mapsto P^\beta: (s, t) \mapsto P_{s \rightarrow t}^\beta = \begin{cases} \mathbf{P}^{\beta|[\sigma, t]} & \text{for } s \leq t \\ (\mathbf{P}^{\beta|[\sigma, s]})^{-1} & \text{for } s \geq t \end{cases} \quad \beta: J \rightarrow M \quad s, t \in J. \quad (5.6)$$

Proof. Suppose I satisfies (4.2). To begin with, we notice that (4.2) imply (3.4) and (3.5) by virtue of proposition 4.1. Equation (2.6) follows from (5.1), and (3.5). Equation (2.7) is a consequence of (5.1), (3.5) and (3.6) for $\gamma: [0, 1] \rightarrow M$ and $\tau_-(t) = 1 - t$ for $t \in [0, 1]$:

$$\mathbf{l}^\gamma = I_{0 \rightarrow 1}^{\gamma \circ \tau_-} = I_{\tau_-(0) \rightarrow \tau_-(1)}^\gamma = I_{1 \rightarrow 0}^\gamma = (I_{0 \rightarrow 1}^\gamma)^{-1} = (\mathbf{l}^\gamma)^{-1}.$$

To prove (2.8), we set $\gamma_{1,2}: [0, 1] \rightarrow M$, $\tau_1(t) = 2t$ for $t \in [0, 1/2]$ and $\tau_2(t) = 2t - 1$ for $t \in [1/2, 1]$ and, applying equations (5.1), (3.4), (3.5), (3.6) and (3.2), we find:

$$\begin{aligned} \mathbf{l}^{\gamma_1 \gamma_2} &= I_{0 \rightarrow 1}^{\gamma_1 \gamma_2} = I_{1/2 \rightarrow 1}^{\gamma_1 \gamma_2} \circ I_{0 \rightarrow 1/2}^{\gamma_1 \gamma_2} = I_{1/2 \rightarrow 1}^{\gamma_1 \gamma_2| [1/2, 1]} \circ I_{0 \rightarrow 1/2}^{\gamma_1 \gamma_2| [0, 1/2]} \\ &= I_{1/2 \rightarrow 1}^{\gamma_2 \circ \tau_2} \circ I_{0 \rightarrow 1/2}^{\gamma_1 \circ \tau_1} = I_{\tau_2(1/2) \rightarrow \tau_2(1)}^{\gamma_2} \circ I_{\tau_1(0) \rightarrow \tau_1(1/2)}^{\gamma_1} = I_{0 \rightarrow 1}^{\gamma_2} \circ I_{0 \rightarrow 1}^{\gamma_1} = \mathbf{l}^{\gamma_2} \circ \mathbf{l}^{\gamma_1}. \end{aligned}$$

^e In particular, one can set $\sigma = \min(s, t)$ and $\tau = \max(s, t)$ or, if J is a closed interval, define σ and τ as the end points of J , i.e. $J = [\sigma, \tau]$.

Next, equation (2.9) is a direct corollary from (5.1) and (3.3). At last, the properties (v) and (vi) are trivial consequences from (5.1) and respectively (4.2a)-(4.2b) and (4.2c). So, $\mathbf{l}$ is a parallel transport, which, evidently, satisfies (2.13) with $\mathbf{l}$ for $\mathbf{P}$, if (5.2) is valid.

Conversely, suppose $\mathbf{P}$ is a parallel transport and P is defined via (5.4), or, equivalently (see remark 5.1), via (5.6). The verification of (3.1)–(3.3) can be done by applying (5.3) and (2.6)–(2.9). Therefore P is a parallel transport along paths. Similarly, the conditions (4.2) follow from the properties (v) and (vi) of the parallel transport $\mathbf{P}$. At last, the validity of (2.9), with P for I , is a consequence from (2.13) and (5.6). $\square$

Definition 5.1. A transport I along paths which has the properties (4.2) will be called *parallel transport along paths*.

Theorem 5.1 simply says that there is a bijective correspondence between the parallel transports along paths and the parallel transports.

Definition 5.2. If I is a parallel transport along paths, then we say that the parallel transport (5.1) is *generated by* (defined by, assigned to) I . Respectively, if $\mathbf{P}$ is a parallel transport, then we say that the (parallel) transport along paths (5.4) is *generated by* (defined by, assigned to) $\mathbf{P}$.

Corollary 5.1. The parallel transport $\mathbf{l}$ generated by the connection Δ^I , assigned to a transport I along paths, is a parallel transport, i.e. it satisfies (2.6)–(2.12) with $\mathbf{l}$ for $\mathbf{P}$.

Proof. This result is a particular case of proposition 2.1. An alternative proof can be carried out by using (3.1)–(4.2), (4.10), and the definitions of inverse path and product of paths. The assertion is also a consequence of (4.10) and theorem 5.1. $\square$

Proposition 5.1. If I is a parallel transport along paths and $\mathbf{l}$ is the assigned to it parallel transport, then the parallel transport along paths defined by $\mathbf{l}$ coincides with I . Conversely, if $\mathbf{P}$ is a parallel transport and P is the parallel transport along paths generated by $\mathbf{P}$, then the parallel transport defined by P coincides with $\mathbf{P}$.

Proof. The assertions are consequences from definition 5.2, theorem 5.1 and remark 5.1. $\square$

6. Relations between connections and (parallel) transports along paths

Until this point, we have studied how a transport along paths generates a connection (theorem 4.1) and parallel transport (proposition 4.3). Besides, theorem 5.1 establishes a bijective correspondence between particular class of transports along paths and mappings having (some of) the main properties of the parallel transports generated by connections. Below we shall pay attention, in a sense, to the opposite links, starting from a connection on a bundle.

Proposition 6.1. Let $\mathbf{P}$ be the parallel transport assigned to a connection Δ^h on a bundle (E, π, M) . The mapping

$$P: \gamma \mapsto P^\gamma: (s, t) \mapsto P_{s \rightarrow t}^\gamma \quad \gamma: J \rightarrow M \quad (6.1a)$$

defined by

$$P_{s \rightarrow t}^\gamma = \begin{cases} \mathbf{P}^{\gamma|_{[s, t]}} & \text{for } s \leq t \\ (\mathbf{P}^{\gamma|_{[t, s]}})^{-1} & \text{for } s \geq t \end{cases} \quad (6.1b)$$

is a transport along paths in (E, π, M) . Moreover, P is parallel transport along paths, i.e. it satisfies the equations (4.2) from theorem 4.1 with P for I .

Proof. One should check the conditions (3.1)–(3.3) and (4.2) with P for I . The relations (3.1)–(3.3) follow directly from definition 2.2 of a parallel transport generated by a connection. The rest conditions are consequences of (6.1) and a simple, but tedious, application of the properties (2.6)–(2.8) of the parallel transports. Alternatively, this proposition is a consequence of the second part of theorem 5.1 and remark 5.1. $\square$

Remark 6.1. Applying (2.6)–(2.8), the reader can verify that

$$P_{s \rightarrow t}^\gamma = F^{-1}(t; \gamma) \circ F(s; \gamma) \quad (6.2)$$

with

$$F(r; \gamma) = \begin{cases} P^{\gamma| [r, w]} & \text{for } r \leq w \\ (P^{\gamma| [w, r]})^{-1} & \text{for } r \geq w \end{cases} \quad r = s, t \quad (6.3)$$

for any (arbitrarily) fixed $w \in J$. This result is a special case of the general structure of the transports along paths [41, theorem 3.1].

Definition 6.1. The parallel transport along paths, defined by a connection Δ^h on a bundle through proposition 6.1, will be called parallel transport along paths *assigned to* (defined by, generated by) the connection Δ^h .

Corollary 6.1. Let Δ^I be the connection generated by a parallel transport I along paths according to theorem 4.1. If l is the parallel transport assigned to Δ^I , then the transport along paths assigned to l (or Δ^I), as described in proposition 6.1, coincides with the initial transport I along paths.

Proof. Substitute (4.10) into (6.1), with I for P and l for P . $\square$

Corollary 6.2. Let P be the transport along paths assigned to a connection Δ^h (via its parallel transport P) according to proposition 6.1. The connection Δ^P generated by P , as described in theorem 4.1, coincides with the initial connection Δ^h , $\Delta^P = \Delta^h$.

Proof. On one hand, if $p \in E$, the space Δ_p^P consists of the vectors tangent at s_0 to the paths $\tilde{\gamma}_{s_0, p}: t \mapsto P_{s_0 \rightarrow t}^\gamma(p)$, with $\gamma: J \rightarrow M$, $s_0 \in J$, and $\pi(p) = \gamma(s_0)$, due to theorem 4.1. On another hand, Δ_p^h consists of the vectors tangent at s_0 to the paths $\tilde{\gamma}_{s_0, p}: t \mapsto P^{\gamma| [s_0, t]}(p)$, by virtue of definition 2.2. Equation (6.1b) says that both types of paths coincide, $\tilde{\gamma}_{s_0, p} = \tilde{\gamma}_{s_0, p}$, so that their tangent vectors at $t = s_0$ are identical and, consequently, Δ_p^P and Δ_p^h are equal as sets, $\Delta_p^P = \Delta_p^h$, for all $p \in E$. $\square$

7. Recapitulation

Roughly speaking, the above series of results says that a connection Δ^h is equivalent to a mapping P (the assigned to it parallel transport) satisfying (2.5)–(2.12) or to a mapping P (the assigned to it parallel transport along paths) satisfying (3.1)–(3.3) and (4.2) (with P for I). Besides, the smoothness of Δ^h is equivalent to the one of P or P . Let us summarize these results as follows.

Theorem 7.1. Given a connection Δ^h on a bundle (E, π, M) , there exists a unique parallel transport I along paths in (E, π, M) which generates Δ^h via (4.3), i.e. $\Delta^I = \Delta^h$. Besides, the parallel transport P defined by Δ^h is given by (5.1), i.e. $P = \mathsf{l}$.

Proof. See theorem 4.1, proposition 4.3 and theorem 5.1. $\square$

Theorem 7.2. *Given a parallel transport I along paths in a bundle (E, π, M) , then there exists a unique connection Δ^h on (E, π, M) such that the parallel transport P along paths assigned to Δ^h coincides with I , $P = I$. Besides, the connection Δ^I generated by I is identical with Δ^h , $\Delta^I = \Delta^h$.*

Proof. Apply theorem 4.1 and corollaries 6.1 and 6.2. □

Theorem 7.3. *Given a parallel transport along paths in a bundle, there is a unique (axiomatically defined) parallel transport generating it. Conversely, given a parallel transport, there is a unique parallel transport along paths generating it.*

Proof. This statement is a reformulation of theorem 5.1. □

Theorem 7.4. *Given a parallel transport P , there exists a unique connection Δ^h generating it. Besides, the parallel transport assigned to Δ^h coincides with P .*


Proof. See theorems 7.3 and 7.2 and definitions 2.2 and 5.2. □

Theorem 7.5. *Given a connection Δ^h , there is a unique parallel transport P such that the defined by it parallel transport P along paths generates Δ^h , $\Delta^P = \Delta^h$. Besides, P coincides with the parallel transport assigned to Δ^h .*

Proof. Apply theorems 7.1 and 7.3. □

We end with the main moral of the We can summarize the above consiabove theorems. In a case of a differentiable bundle, the concepts “connection,” “(axiomatically defined) parallel transport” and “parallel transport along path” are equivalent in a sense that there are bijective mappings between the sets of these objects; thus, we have the commutative diagram shown on figure 7.1 in which the mappings indicated are described via theorems 7.1–7.5.

Figure 7.1: Mappings between the sets of parallel transports, connections and parallel transports along paths in differentiable bundles



Besides, if one of these objects is smooth, so are the other ones corresponding to it via the bijections constructed in the present work.

8. The case of topological bundles

Now, following some ideas of [45], we want to say a few words concerning bundles which are *not* differentiable. The definition 2.1 of a connection is strongly related to the concept of the tangent bundle $T(E)$ of the bundle space E and, consequently, it is senseless in a case

when E is a manifold of class C^0 of a topological space. As we have demonstrated above (see theorems 7.1 and 7.2), a connection can equivalently be described via a suitable parallel transport which, by definition 2.3, is a mapping possessing the properties (2.6)–(2.12). One observes at first sight that the properties (2.6)–(2.9) do not depend on the differentiable structure of the bundle and, on the contrary, the properties (2.10)–(2.12) are senseless if E is not of class C^1 or higher. This separation of the properties of the parallel transports assigned to connections to ones that depend and do not depend on the differentiable structure of the bundles points to a natural way for generalizing the concept of a parallel transport (and hence of a connection) on bundles with class of smoothness not higher than C^0 .

Definition 8.1. A parallel transport P in a topological bundle (E, π, B) is a mapping (2.5) (with B for M) which has the properties (2.6)–(2.9). Such a mapping will also be called *topological parallel transport* if there is a risk to mix it with the one defined via definition 2.3 in a case of a differentiable bundle.

Proposition 8.1. A parallel transport in a differentiable bundle of class C^1 is a topological parallel transport.

Proof. Compare definitions 2.3 and 8.1. $\square$

Relying on our experience with connections and parallel transports in differentiable bundles, the concepts “connection” and “(topological) transport” in topological bundles should be identified and, hence considered as equivalent. As the following result shows, the set of (topological) parallel transports is in a bijective correspondence with certain subset of transports along paths in topological bundles.

Theorem 8.1. Let I be a transport along paths in an arbitrary topological bundle (E, π, B) and $\gamma: [\sigma, \tau] \rightarrow B$. If I satisfies the conditions (3.4) and (3.5), then the mapping

$$l: \gamma \mapsto l^\gamma := I_{\sigma \rightarrow \tau}^\gamma: \pi^{-1}(\gamma(\sigma)) \rightarrow \pi^{-1}(\gamma(\tau)) \quad \gamma: [\sigma, \tau] \rightarrow M \quad (8.1)$$

is a parallel transport, i.e. it possess the properties (2.6)–(2.9), with l for P .

Conversely, suppose the mapping

$$P: \gamma \mapsto P^\gamma: \pi^{-1}(\gamma(\sigma)) \rightarrow \pi^{-1}(\gamma(\tau)) \quad \gamma: [\sigma, \tau] \rightarrow M \quad (8.2)$$

is a parallel transport, i.e. satisfies (2.6)–(2.9), and define the mapping

$$P: \beta \mapsto P^\beta: (s, t) \mapsto P_{s \rightarrow t}^\beta = \begin{cases} P^{\beta|_{[s, t]}} & \text{for } s \leq t \\ (P^{\beta|_{[t, s]}})^{-1} & \text{for } s \geq t \end{cases} \quad \beta: J \rightarrow B \quad s, t \in J. \quad (8.3)$$

Then the mapping (8.3) is a transport along paths in (E, π, B) , which transport satisfies the conditions (3.4) and (3.5), with P for I .

Proof. To prove the first part of the theorem, we define $\tau_-: t \mapsto 1 - t$, $\tau_1: t \mapsto 2t$ and $\tau_2: t \mapsto 2t - 1$ for $t \in [0, 1]$. Applying (3.2)–(3.5) and using the notation of proposition 2.1, we get:

$$\begin{aligned} l^{\gamma \circ \chi} &= I_{\sigma \rightarrow \tau}^{\gamma \circ \chi} = I_{\chi(\sigma) \rightarrow \chi(\tau)}^\gamma = l^\gamma \\ l^{\gamma^-} &= I_{0 \rightarrow 1}^{\gamma \circ \tau_-} = I_{\tau_-(0) \rightarrow \tau_-(1)}^\gamma = I_{1 \rightarrow 0}^\gamma = (I_{0 \rightarrow 1}^\gamma)^{-1} = (l^\gamma)^{-1} \\ l^{\gamma_1 \gamma_2} &= I_{0 \rightarrow 1}^{\gamma_1 \gamma_2} = I_{1/2 \rightarrow 1}^{(\gamma_1 \gamma_2)|_{[1/2, 1]}} \circ I_{0 \rightarrow 1/2}^{(\gamma_1 \gamma_2)|_{[0, 1/2]}} = I_{1/2 \rightarrow 1}^{\gamma_2 \circ \tau_2} \circ I_{0 \rightarrow 1/2}^{\gamma_1 \circ \tau_1} = I_{0 \rightarrow 1}^{\gamma_2} \circ I_{0 \rightarrow 1}^{\gamma_1} = l^{\gamma_2} \circ l^{\gamma_1} \\ l^{\gamma_{r, x}} &= I_{r \rightarrow r}^{\gamma_{r, x}} = \text{id}_{\pi^{-1}(x)}. \end{aligned}$$

To prove the second part, we first note that (3.3) and (3.5) follow from (2.9) and (2.6), respectively, due to (8.3). As a result of (8.3), we have, e.g. for $s \leq t$,

$$P_{s \rightarrow t}^{\gamma|J'} = \mathbf{P}^{(\gamma|J')|[s,t]} = \mathbf{P}^{\gamma|[s,t]} = P_{s \rightarrow t}^{\gamma}$$

by virtue of $[s, t] \subseteq J'$. At last, we shall prove (3.2) for $r \leq s \leq t$; the other cases can be proved similarly. Let us fix some orientation-preserving bijections $\tau': [0, 1] \rightarrow [r, s]$ and $\tau'': [0, 1] \rightarrow [s, t]$. Then:

$$\begin{aligned} P_{s \rightarrow t}^{\beta} \circ P_{r \rightarrow s}^{\beta} &= \mathbf{P}^{\beta|[s,t]} \circ \mathbf{P}^{\beta|[r,s]} = \mathbf{P}^{\beta \circ \tau''} \circ \mathbf{P}^{\beta \circ \tau'} = \mathbf{P}^{(\beta \circ \tau')(\beta \circ \tau'')} \\ &= \mathbf{P}^{\beta \circ \tau} = \mathbf{P}^{\beta|[r,t]} = P_{r \rightarrow t}^{\beta} \end{aligned}$$

where $\tau: a \mapsto \tau'(2a)$ for $a \in [0, 1/2]$ and $\tau: a \mapsto \tau''(2a - 1)$ for $a \in [1/2, 1]$, so that $\tau(0) = \tau'(0) = r$ and $\tau(1) = \tau''(1) = t$. $\square$

Remark 8.1. In fact, this proof is contained implicitly in the proof of theorem 5.1.


Definition 8.2. In a topological bundle, a transport I along paths which has the properties (3.4) and (3.5) will be called *parallel transport along paths*. If I is a parallel transport along paths, then we say that the parallel transport (8.1) is *generated by (defined by, assigned to)* I . Respectively, if $\mathbf{P}$ is a parallel transport, then we say that the (parallel) transport along paths (8.3) is *generated by (defined by, assigned to)* $\mathbf{P}$.

Theorem 8.1 simply says that there is a bijective correspondence between the set of parallel transports along paths and the one of parallel transports.

Theorem 8.1 and definition 8.2 are analogues of respectively Theorem 5.1 and definition 5.1 in a case of a topological bundle and contain them as special cases when differentiable bundles are considered. The last assertion is confirmed by the result of exercise 4.1 that equations (3.4) and (3.5) are consequences from the equations (4.2). Thus, in terms of transports along paths, equations (3.4) and (3.5) single out the parallel transports which are independent of the differentiable structure of the bundles in which they act. The same result is expressed by the properties (2.6)–(2.9) of the parallel transports assigned to connections while (2.10)–(2.12) are sensible only in differentiable bundles and single out from the topological parallel transports the ones defined by connections.

We can summarize the above considerations in the commutative diagram presented on figure 8.2 the mappings in which are described via theorem 8.1.

Figure 8.2: Mappings between the sets of (topological) parallel transports and parallel transports along paths in topological bundles



As a conclusion, we can say that in topological bundles the concept of connection does not survive and there is a bijection between the sets of “parallel transports” and “parallel transports along paths” and, in that sense, these concepts are equivalent.

9. The case of vector bundles

The purpose of this section is to be presented some links between transports and connections in vector bundles^f in a case these objects are linear, i.e. when they are compatible with the vector structure of the bundle.

Definition 9.1. A *linear* transport L along paths in $\mathbb{K} = \mathbb{R}, \mathbb{C}$ vector bundle (E, π, M) is a transport L along paths such that

$$L_{s \rightarrow t}^\gamma(\lambda u + \mu v) = \lambda L_{s \rightarrow t}^\gamma(u) + \mu L_{s \rightarrow t}^\gamma(v) \quad \lambda, \mu \in \mathbb{K}, \quad u, v \in \pi^{-1}(\gamma(s)) \quad (9.1)$$

for all paths $\gamma: J \rightarrow M$ and $s, t \in J$.

Definition 9.2 (see [46, p. 354]). A connection on a vector bundle is called *linear* if the assigned to it parallel transport P is a linear mapping along every path in the base space, i.e. for all paths $\gamma: [\sigma, \tau] \rightarrow M$, we have

$$P^\gamma(\rho X) = \rho P^\gamma(X) \quad (9.2a)$$

$$P^\gamma(X + Y) = P^\gamma(X) + P^\gamma(Y), \quad (9.2b)$$

where $\rho \in \mathbb{K}$ and $X, Y \in \pi^{-1}(\gamma(\sigma))$.

Before proceeding with the topic of this section, we shall present some results concening linear transports and linear connections.

If $\{e_a\}$, $a, b = 1, \dots, \dim \pi^{-1}(x)$, $x \in M$, is a frame in E , i.e. $e_a: M \ni x \rightarrow e_a(x)$ with $\{e_a(x)\}$ being a basis in $\pi^{-1}(x)$, then the matrix $\mathbf{L} = [L_b^a]$ of L is defined by $L_{s \rightarrow t}^\gamma(e_a(\gamma(s))) = L_a^b(t, s; \gamma) e_b(\gamma(t))$ and its general form is [46, chapter IV, proposition 3.4]

$$\mathbf{L}(t, s; \gamma) = F^{-1}(t; \gamma) F(s; \gamma) \quad (9.3)$$

for some non-degenerate matrix-valued function F . The coefficients $\Gamma_b^a(s, \gamma)$ of a C^1 linear transport L are defined by

$$\mathbf{\Gamma}(s, \gamma) := [\Gamma_b^a(s, \gamma)] := \left. \frac{\partial \mathbf{L}(s, t; \gamma)}{\partial t} \right|_{t=s} = F^{-1}(s; \gamma) \frac{dF(s; \gamma)}{ds} \quad (9.4)$$

and determine the derivation along paths corresponding to L [46, chapter IV, section 3.3].

If $\{X_I\}$, $I, J = 1, \dots, \dim M, \dots, \dim M + \dim \pi^{-1}(x)$, $x \in M$, is a frame in $T(E)$ adapted to $\{\frac{\partial}{\partial u^I}\}$ (or to $\{u^I\}$), with $\{u^I\}$ being vector bundle coordinates in E , for a connection Δ^h , then [46, p. 368]

$$X_\mu = \frac{\partial}{\partial u^\mu} + \Gamma_\mu^\beta \frac{\partial}{\partial u^\beta} \quad X_a = \frac{\partial}{\partial u^a} \quad (9.5)$$

where $\mu, \nu = 1, \dots, \dim M$ and Γ_μ^b are the (2-index) coefficients of Δ^h . According to theorem 4.1 in [46, p. 354], the connection Δ^h is linear iff

$$\Gamma_\mu^a(p) = -\Gamma_{b\mu}^a(\pi(p)) u^b(p) = -((\Gamma_{b\mu}^a \circ \pi) \cdot u^b)(p) \quad p \in E, \quad (9.6)$$

for some functions $\Gamma_{b\mu}^a$ known as the (3-index) coefficients of Δ^h .

Now we shall present some important results.

Proposition 9.1. *In a case of a vector bundle:*

- (a) *A parallel transport I along paths is linear if and only if the assigned to it parallel transport $\mathbf{l}$ is linear;*
- (b) *A parallel transport P is linear if and only if the generated by it parallel transport P along paths is linear.*

^f Details on the vector bundle theory can be found, for instance, in [32].

Proof. The results follow from theorem 5.1 and proposition 5.1. $\square$

Proposition 9.2. *In a case of a vector bundle:*

(a) *The linear connections are the only connections that can be generated by linear parallel transports along paths;*

(b) *The linear parallel transports along paths are the only transports along paths that can be generated by linear connections (via the assigned to them parallel transports).*

Proof. The assertions are consequences from definition 9.1 and propositions 9.1 and 6.1. $\square$

Theorem 9.1. *Let Δ^h be a linear connection on a vector bundle (E, π, M) , E and M being C^1 manifolds. Let $\{u^I\} = \{x^\mu \circ \pi, u^a\}$, with $\{x^\mu\}$ being a coordinate system on M , be vector bundle coordinates on E generated by a frame $\{e_a\}$ in E , $\{X_I\}$ be the frame adapted to $\{u^I\}$ and $\Gamma_{b\mu}^a$ be the 3-index coefficients of Δ^h in $\{X_I\}$ (see equation (9.6)). If P is the parallel transport along paths assigned to Δ^h , then the coefficients of P in $\{e_a\}$ along a C^1 path $\gamma: J \rightarrow M$ are ($s \in J$)*

$$\Gamma_b^a(s; \gamma) = \Gamma_{b\mu}^a(\gamma(s)) \dot{\gamma}^\mu(s). \quad (9.7)$$

Conversely, if I is a C^1 linear transport along paths in (E, π, M) and in $\{e_a\}$ its coefficients have a representation (9.7) along every C^1 path γ for some functions $\Gamma_{b\mu}^a$, then there exists a unique linear connection Δ^h on (E, π, M) such that the parallel transport P along paths generated by Δ^h coincides with I , $P = I$; besides, $\Gamma_{b\mu}^a$ are exactly the 3-index coefficients of Δ^h .

Proof. According to corollaries 6.1 and 6.2 and proposition 4.2, the Δ^h -horizontal lift of γ through $p \in \pi^{-1}(\gamma(s_0))$, $s_0 \in J$ is given by $\bar{\gamma}_{s_0,p}(s) = P_{s_0 \rightarrow s}^\gamma(p)$. Then, in a (local) frame $\{e_a\}$ generating vector bundle coordinates $\{u^I\} = \{u^\mu = x^\mu \circ \pi, u^a\}$, $\{x^\mu\}$ being a coordinate system on M , we have

$$\bar{\gamma}_{s_0,p}(t) = P_b^a(t, s_0; \gamma) p^b e_a(\gamma(t))$$

due to the definition of the matrix elements P_b^a of P . Using (9.3) and (9.4), we find from here $\dot{\gamma}_{s_0,p}^\mu(t) = \dot{\gamma}^\mu(t)$, which is other form of lemma 4.1, and

$$\dot{\gamma}_{s_0,p}^a(s) = -\Gamma_b^a(s; \gamma) \bar{\gamma}_{s_0,p}^b(s). \quad (9.8)$$

On another hand, from the parallel transport equation [46, p. 349, eq. (3.39'a)], we get

$$\dot{\gamma}_{s_0,p}^a(s) = \Gamma_\mu^a(\bar{\gamma}_{s_0,p}(s)) \dot{\gamma}_{s_0,p}^\mu(s),$$

where Γ_μ^a are the 2-index coefficients of Δ^h . Inserting here (9.6) and comparing the result with the previous displayed equation, we obtain (9.7), due to $\pi \circ \bar{\gamma}_{s_0,p} = \gamma$ and (4.5).

Conversely, suppose (9.7) holds. Then I satisfies the conditions (a)–(c) from theorem 4.1, so (4.3) defines a connection Δ^h . Then the parallel transport P along paths generated by Δ^h coincides with I , $P = I$, by virtue of corollary 6.1. As a result of proposition 4.2, (9.7) and (9.8), the Δ^h horizontal lift of γ is such that

$$\dot{\gamma}_{s_0,p}(s) = -\Gamma_{b\mu}^a(\gamma(s)) \dot{\gamma}^\mu(s) \bar{\gamma}_{s_0,p}^b(s).$$

Comparing this result with the parallel transport equation [46, p. 349, eq. (3.39'a)], we get (see also (4.5))

$$\Gamma_\mu^a(\bar{\gamma}_{s_0,p}(s)) = -\Gamma_{b\mu}^a(\gamma(s)) \bar{\gamma}_{s_0,p}^b(s)$$

which entails $\Gamma_\mu^a = -(\Gamma_{b\mu}^a \circ \pi) \cdot u^b$ due to $\pi \circ \bar{\gamma}_{s_0,p} = \gamma$ and the arbitrariness of γ and $p \in \pi^{-1}(\gamma(s_0))$. At last, according to the above-cited theorem 4.1 in [46, p. 349] the connection Δ^h is linear. $\square$

Theorem 9.1 simply means that a linear transport along paths is a parallel transport along paths (generated by a linear connection) if and only if equation (9.7) holds in some (and hence in any) frame $\{e_a\}$. In this sense, a linear transport along paths is equivalent to a linear connection iff equation (9.7) is valid in some (and hence in any) frame $\{e_a\}$. We can also say that theorem 9.1 describes the subset of all linear transports along paths that can be generated by linear connections, viz. these are the transports with coefficients given by equation (9.7).

As an application of theorem 9.1, we shall prove that the evolution transport arising in bundle quantum mechanics [47] is generally *not a parallel* linear transport along paths.^g To show this, we note that the coefficients of the evolution transport are [48, eq. (2.22)]

$$\Gamma_b^a(t; \gamma) = -\frac{1}{i\hbar}(\mathbf{H}_\gamma^m(t))_b^a \quad (9.9)$$

where i is the imaginary unit, $\hbar$ is the Planck's constant (divided by 2π) and $\mathbf{H}_\gamma^m(t)$ is the matrix-bundle Hamiltonian. One can choose a special frame which is independent of γ and is such that [48, remark 2.1] $\mathbf{H}_\gamma^m(t) = \mathcal{H}(t)$, where $\mathcal{H}(t)$ is the matrix of the usual Hilbert space Hamiltonian. So, in this frame, we have

$$\Gamma_b^a(t; \gamma) = -\frac{1}{i\hbar}\mathcal{H}_b^a(t) \quad (9.10)$$

with $\mathcal{H}_b^a(t)$ being the matrix elements of the system Hamiltonian which may depend on the (time) parameter t but does not depend on the path γ . Therefore equation (9.7) is generally not valid and consequently the evolution transport cannot be generated by a linear connection in the general case. In particular, this conclusion is valid for all time-independent Hamiltonians, $\frac{\partial \mathcal{H}(t)}{\partial t} = 0$.

10. Conclusion

The concept “parallel transport” precedes historically the one of a “connection” and was first clearly formulated in the work [33] of Levi Civita on a parallel transport of a vector in Riemannian geometry.^h The connection theory was formulated approximately during the period 1920–1949 in a series of works on particular connections and their subsequent generalizations and has obtained an almost complete form in 1950–1955 together with the clear formulation of the concepts “manifold”ⁱ and “fibre bundle”^j [26]. It should be noted the contribution of Elie Cartan in the theory of linear connections [52, 53]. Connections as splitting of the second tangent bundle $T(T(M))$ over a manifold M into vertical and horizontal parts were first introduced by Ehresmann in [54] and sometimes are referred as “Ehresmann connections”. The concept of connection on general fibre bundle was established at about 1970 [55]. During that period, with a few exceptional works, the ‘parallel transport’ was considered as a secondary concept, defined by means of the one of a ‘connection’. Later, as we pointed at the beginning of Sect. 3, there appear several attempts for axiomatizing the concept of a parallel transport and by its means the connection theory to be constructed; e.g., this approach is developed deeply in vector bundles in [32].

The main merit from the present paper is that now we have a necessary and sufficient conditions when in differentiable bundles an axiomatically defined parallel transport (or a

^g The below-presented proof is not quite rigorous as theorem 9.1 is proved in the finite-dimensional case while the evolution transport acts in generally infinite dimensional Hilbert bundle.

^h See also the early work [49].

ⁱ The theory of manifolds was worked out by H. Whitney during the 1930s [50] and [51] seems to be one of the first its systematic expositions.

^j The first edition of the book [8] is the first detailed and complete presentation of fibre bundle theory. Later the book [11] became a modern standard for references on bundle theory.

parallel transport along paths) defines a unique connection (with suitable properties) and *vice versa*. Moreover, the concepts “connection,” “(axiomatically defined) parallel transport,” and “parallel transport along path” are equivalent in a sense that there are bijective mappings between the sets of these objects. In topological bundles, we have established a bijection between the sets of parallel transports and of parallel transports along paths. However, the concept of a parallel transport admits a generalization to the one of a transport along paths and there exist transports along paths that cannot be generated by connections or parallel transports. In vector bundles, we have found a condition ensuring an equivalence between linear transports and linear connections.

Acknowledgments

This work was partially supported by the National Science Fund of Bulgaria under Grant No. F 1515/2005.

The author thanks Professor Phillip E. Parker (Department of Mathematics and Statistics, Wichita State University Wichita, Kansas, USA) for some historical comments on connection theory.

References

- [1] S. Kobayashi. Theory of connections. *Ann. mat. pure ed appl.*, 43:119–194, 1957.
- [2] S. Kobayashi and K. Nomizu. *Foundations of Differential Geometry*, volume I. Interscience Publishers, New York-London, 1963. Russian translation: Nauka, Moscow, 1981.
- [3] R. K. Sachs and H. Wu. *General Relativity for Mathematicians*. Springer-Verlag, New York-Heidelberg-Berlin, 1977.
- [4] C. Nash and S. Sen. *Topology and Geometry for physicists*. Academic Press, London-New York, 1983.
- [5] F. W. Warner. *Foundations of differentiable manifolds and Lie groups*. Springer-Verlag, New York-Berlin-Heidelberg-Tokyo, 1983. Russian translation: Mir, Moscow, 1987.
- [6] R. L. Bishop and R. J. Crittenden. *Geometry of Manifolds*. Academic Press, New York-London, 1964.
- [7] K. Yano and M. Kon. *Structures on Manifolds*, volume 3 of *Series in Pure Mathematics*. World Scientific Publ. Co., Singapore, 1984.
- [8] Norman Steenrod. *The topology of fibre bundles*. Princeton University Press, Princeton, ninth edition, 1974. First edition: Princeton University Press, 1951.
- [9] R. Sulanke and P. Wintgen. *Differential geometry and fibre bundles*. Mir, Moscow, 1975. In Russian. Translation from the German original: Differentialgeometrie und Faserbündel, VEB Deutscher Verlag der Wissenschaften, Berlin, 1972.
- [10] Y. Choquet-Bruhat et al. *Analysis, manifolds and physics*. North-Holland Publ. Co., Amsterdam, 1982.
- [11] D. Husemoller. *Fibre bundles*. McGraw-Hill Book Co., New York-St. Louis-San Francisco-Toronto-London-Sydney, 1966. Other editions: Springer Verlag, 1975, 1994, 1998. Russian translation: Mir, Moscow, 1970.

- [12] A. S. Mishchenko. *Vector fibre bundles and their applications*. Nauka, Moscow, 1984. In Russian.
- [13] R. Hermann. *Vector bundles in mathematical physics*, volume I. W. A. Benjamin, Inc., New York, 1970.
- [14] W. Greub, S. Halperin, and R. Vanstone. *De Rham cohomology of manifolds and vector bundles*, volume 1 of *Connections, Curvature, and Cohomology*. Academic Press, New York and London, 1972.
- [15] M. F. Atiyah. *K-theory*. Harvard Univ, Cambridge, Mass., 1965.
- [16] R. Dandoloff and W. J. Zakrzewski. Parallel transport along a space curve and related phases. *J. Phys. A: Math. Gen.*, 22:L461–L466, 1989.
- [17] I. Tamura. *Topology of foliations*. Mir, Moscow, 1979. Russian translation from 1976 Japanese original.
- [18] N. J. Hicks. *Notes on differential geometry*. D. Van Nostrand Comp., Inc., Princeton, 1965.
- [19] S. Sternberg. *Lectures on differential geometry*. Chelsea Publ. Co., New York, 1983. First edition: Prentice Hall, Inc. Englewood Cliffs, N.J., 1964. Russian translation: Mir, Moscow, 1970.
- [20] Maïdo Rahula. *New problems in differential geometry*, volume 8 of *Series on Soviet and East European Mathematics*. World Scientific, Singapore-New Jersey-London-Hong Kong, 1993. 172 p.
- [21] L. Mangiarotti and G. Sardanashvily. *Connections in classical and quantum field theory*. World Scientific, Singapore-New Jersey-London-Hong Kong, 2000.
- [22] Ülo G. Lumiste. To the foundations of the global connection theory. In *Scientific writings of the Tartu state university*, number 150 in Works on mathematics and mechanics, IV, pages 69–107. University of Tartu, Tartu, 1964. In Russian.
- [23] Ülo G. Lumiste. Homogeneous fibre bundles with connection and their immersions. In *Works of the Geometrical seminar*, volume 1, pages 191–237. VINITI, Moscow, 1966. In Russian.
- [24] C. Teleman. *Elements of topology and differentiable manifolds*. Mir, Moscow, 1967. In Russian. Translation from the Rumanian original: Elemente de topologie si varietăți diferentiabile, București, 1964. German translation: Grundzüge der Topologie und differenzierbare Mannigfaltigkeiten (Autorisierte und vom Autor erg. Übersetzung aus dem Rumänischen: Horst Antelmann.) Mit 22 Abbildungen, Berlin, Deutscher Verlag der Wissenschaften, 1968, 411 p.
- [25] Peter Dombrowski. Krümmungsgrößen gleichungsdefinierter Untermannigfaltigkeiten Riemannscher Mannigfaltigkeiten. *Mathematische Nachrichten*, 38(3/4):133–180, 1968.
- [26] Ülo G. Lumiste. Connection theory in fibre bundles. In *Science review*, Mathematics: Algebra. Topology. Geometry. 1969, pages 123–168. VINITI, Moscow, 1971. In Russian.
- [27] *Mathematical encyclopedia*, chief editor Vinogradov I. M., volume 4, Moscow, 1984. Sovetskaya Entsiklopediya (Soviet encyclopedia). In Russian. English translation in: Encyclopedia of Mathematics, vol. I–VI, Kluwer, 1995.

- [28] B. Durhuus and J. M. Leinaas. On the loop space formulation of gauge theories. Preprint TH 3110, CERN, Geneva, 1981.
- [29] O. M. Khudaverdian and A. S. Schwarz. A new comments on the string representation of gauge fields. *Phys. Lett. B*, 91(1):107–110, 1980.
- [30] N. M. Ostianu, V. V. Rizhov, and P. I. Shveikin. Article on the scientific investigations of German Fedorovich Laptev. In *Works of the Geometrical seminar*, volume 4, pages 7–70. VINITI, Moscow, 1973. In Russian.
- [31] P. Nikolov. On the correspondence between infinitesimal and integral description of connections. Internal Report, IC/81/196, ICTP, Trieste, 1981.
- [32] Walter A. Poor. *Differential geometric structures*. McGraw-Hill Book Company Inc., New York, 1981.
- [33] T. Levi-Civita. Nazione di parallelismo in una variete qualunque e conseguente spetificazione geometrica della curvaturre riemanniano. *Rend. Palermo*, 42:73–205, 1917. In Italian.
- [34] S. Kobayashi and K. Nomizu. *Foundations of Differential Geometry*, volume I and II. Interscience Publishers, New York-London-Sydney, 1963 and 1969. Russian translation: Nauka, Moscow, 1981.
- [35] Sze-Tsen Hu. *Homotopy Theory*. Academic Press, New York-London, 1959.
- [36] O. Ya. Viro and D. B. Fuks. Introduction to homotopy theory. In *Reviews of science and technique*, volume 24 of sec. “Modern problems in mathematics”, pages 6–121. VINITI, Moscow, 1988. In Russian.
- [37] B. A. Dubrovin, S. P. Novikov, and A. T. Fomenko. *Modern geometry: methods and applications, pt. 1*, volume 93 of *Graduate text in Mathematics: The geometry of surfaces, transformation groups, and fields*. Springer, New York, 2 edition, 1992. Translation from the original Russian ed., Moscow, Nauka, 1979.
- [38] B. A. Dubrovin, S. P. Novikov, and A. T. Fomenko. *Modern geometry: methods and applications, pt. 2*, volume 104 of *Graduate text in Mathematics: The geometry and topology of manifolds*. Springer, New York, 1985. Translation from the original Russian ed., Moscow, Nauka, 1979.
- [39] B. A. Dubrovin, S. P. Novikov, and A. T. Fomenko. *Modern geometry: methods and applications, pt. 3*, volume 124 of *Springer series in Soviet mathematics: Introduction to homotopy theory*. Springer, New York, 1990. Translation from the original Russian ed., Moscow, Nauka, 1984.
- [40] D. Gromoll, W. Klingenberg, and W. Meyer. *Riemannian geometry at large*. Mir, Moscow, 1971. Russian translation from the German original: Riemannsche Geometrie im groen, Springer, Berlin-Heidelberg-New York, 1968.
- [41] Bozhidar Z. Iliev. Transports along paths in fibre bundles. General theory. JINR Communication E5-93-299, Dubna, 1993. 26 pp.
<http://arXiv.org> e-Print archive, E-print No. math.DG/0503005, March 1, 2005.
- [42] Bozhidar Z. Iliev. Transports along paths in fibre bundles. II. Ties with the theory of connections and parallel transports. JINR Communication E5-94-16, Dubna, 1994. 24 pp.
<http://arXiv.org> e-Print archive, E-print No. math.DG/0503006, March 1, 2005.

- [43] Bozhidar Z. Iliev. Normal frames and linear transports along paths in vector bundles. *JP Journal of Geometry and Topology*, 5(3):187–250, 2005.
<http://arXiv.org> e-Print archive, E-print No. gr-qc/9809084, September 1998 (last revision: March 2005).
- [44] Bozhidar Z. Iliev. Normal frames for derivations and linear connections and the equivalence principle. *Journal of Geometry and Physics*, 45(1–2):24–53, February 2003.
 DOI No. 10.1016/S0393-0440(02)00119-5
<http://arXiv.org> e-Print archive, E-print No. hep-th/0110194, October 2001.
- [45] Bozhidar Z. Iliev. On the axiomatization of “parallel transport” and its generalization. *Tensor, N. S.*, 62(2):170–181, 2006.
 Report presented at the 8th International Conference of Tensor Society on “Differential geometry, functional and complex analysis, informatics and their applications”, Varna, Bulgaria, 22 – 26 August, 2005
<http://arXiv.org> e-Print archive, E-print No. math-ph/0510005, October 1, 2005.
- [46] Bozhidar Z. Iliev. *Handbook of Normal Frames and Coordinates*, volume 42 of *Progress in Mathematical Physics*. Birkhäuser, Basel-Boston-Berlin, 2006. 468 pages, Hardcover, ISBN (ISBN-10): 3-7643-7618-X, ISBN-13: 978-3-7643-7618-5, EAN: 9783764376185;
<http://arXiv.org> e-Print archive, E-print No. math.DG/0610037, October 1, 2006.
- [47] Bozhidar Z. Iliev. Fibre bundle formulation of nonrelativistic quantum mechanics. I. Introduction. The evolution transport. *Journal of Physics A: Mathematical and General*, 34(23):4887–4918, 2001. DOI No. 10.1088/0305-4470/34/23/308
<http://arXiv.org> e-Print archive, E-print No. quant-ph/9803084, March 1998.
- [48] Bozhidar Z. Iliev. Fibre bundle formulation of nonrelativistic quantum mechanics. II. Equations of motion and observables. *Journal of Physics A: Mathematical and General*, 34(23):4919–4934, 2001. DOI No. 10.1088/0305-4470/34/23/309
<http://arXiv.org> e-Print archive, E-print No. quant-ph/9804062, April 1998.
- [49] T. Levi-Civita and G. Ricci. Méthodes de calcul différentiel absolu et leurs applications. *Math. Ann.*, B 54:125–201, 1900.
- [50] Hassler Whitney. *Topology: Lecture notes*. Unknown Binding, Harvard, 1963.
- [51] G. de Rham. *Varieties Differentiables*. Hermann, Paris, 1955.
- [52] Elie Cartan. Sur les variétés à connexion projective. *Bulletin de la Société Mathématique*, pages 205–241, 1924.
- [53] E. Cartan. *Geometry of Riemannian spaces*. Math Sci Press, 1983.
- [54] C. Ehresmann. Les connexions infinitésimales dans un espace fibré différentiable. In *Colloque de topologie (espaces fibrés), Bruxelles, 1950.*, pages 29–55. Masson et Cie., Paris, 1951.
- [55] A. Lieberman. Parallélismes. *Journal of differential geometry*, pages 511–519, 1973.